


Account
m and n each is independently an integer of from 0 to 4; p is 1 or 2; and pharmaceutically acceptable salts thereof.

3. A method of blocking or inhibiting tissue factor-dependent activation of factor X and/or factor IX, comprising contacting tissue factor with a TF blocking compound to thereby inhibit formation of a functional complex of factor X or factor IX with tissue factor or TF VIIA, wherein the TF blocking compound is of the following Formula I:



wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O, S, S(O) or S(O₂);

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;
m and n each is independently an integer of from 0 to 4; p is 1 or 2; and pharmaceutically acceptable salts thereof.

In claim 9, line 1, after "claim 7" please delete "or 8".

In claim 10, line 1, after "claim 7" please delete "or 8".

Kindly cancel claim 11.

In claim 12, line 10, please delete "preferably."